



**CINV FOR >6 MONTHS (ABLE TO RECEIVE DEXAMETHASONE BUT NO APREPITANT\*)**

If N&V is not controlled (BARF >4) with recommended antiemetic therapy for chemo, then document BARF below and complete the next checkbox. Consider also anticipatory N&V.

NAME:

CHEMO:

LOW

MODERATE

HIGH

ONDANSETRON

0.15mg/kg STAT  
(Max dose 8mg)

DEXAMETHASONE

No

0.15mg/kg/dose q8h  
(Max dose 8mg)

2mg/dose q12h <0.6m<sup>2</sup>  
4mg/dose q12h >0.6m<sup>2</sup>  
(Max dose 16mg/day max 4 days)

0.15mg/kg/dose q6h<sup>#</sup>  
(Max dose 8mg)

4-6mg/m<sup>2</sup>/dose q12h  
(Max dose 16mg/day max 4 days then review)

SEE RESISTANT CINV

DATE:

BARF SCORE:

- # review frequency after 48 hrs – constipation risk
- **ALL ARE REGULAR ONLY UNTIL 24 HOURS POST CHEMOTHERAPY THEN REVIEW**
- Max dose ASCO 2017

\*Aprepitant is a moderate CYP3A4 inhibitor (definite interactions with dexamethasone; others possible including CPM, Ifos, midaz, oxycodone, anthracyclines)