



CINV FOR >6 MONTHS (ABLE TO RECEIVE BOTH APREPITANT AND DEXAMETHASONE)

If N&V is not controlled (BARF >4) with recommended antiemetic therapy for chemo, then document BARF below and complete the next checkbox. Consider also anticipatory N&V.

NAME:

CHEMO:

LOW

MODERATE

HIGH

ONDANSETRON

0.15mg/kg STAT
(Max dose 8mg)

DEXAMETHASONE

No

APREPITANT[&]

No

0.15mg/kg/dose q8h
(Max dose 8mg)

2mg/dose q12h <0.6m²
4mg/dose q12h >0.6m²
(Max dose 16mg/day max 4 days)

No

0.15mg/kg/dose q6h[#]
(Max dose 8mg)

1-3mg*/m²/dose q12h
(8mg/day as on Aprepitant*)

≥12y or ≥40kg 125mg day 1 and 80mg day 2 and 3
25kg to 40kg 80mg day 1, 2 and 3
15kg to 24kg 80mg day 1 and 40mg day 2 and 3
<15kg 3mg/kg day 1 and 2mg/kg day 2 and 3

SEE
RESISTANT
CINV

DATE:

BARF SCORE:

- * Dose halved already for Aprepitant interaction; max dose from ASCO 2017
- # review frequency after 48hr – constipation risk
- & comes as a 3-day course but may give as single dose (e.g. HD MTX) or for up to 7/7
- **ALL ARE REGULAR ONLY UNTIL 24 HOURS POST CHEMOTHERAPY THEN REVIEW**