



CINV FOR >6 MONTHS (ABLE TO RECEIVE APREPITANT BUT NO DEXAMETHASONE)

If N&V is not controlled (BARF >4) with recommended antiemetic therapy for chemo, then document BARF below and complete the next checkbox. Consider also anticipatory N&V.

NAME:
CHEMO:

LOW

MODERATE

HIGH

ONDANSETRON

0.15mg/kg STAT
(Max dose 8mg)

No

No

METOCLOPRAMIDE

0.15mg/kg/dose q8h
(Max dose 8mg)

0.15mg/kg/dose q8h
(Max 10mg/dose)

No

APREPITANT[&]

0.15mg/kg/dose q6h[#]
(Max dose 8mg)

0.15mg/kg/dose q6-8h
(Max 10mg/dose)

≥12y or ≥40kg 125mg day 1 and 80mg day 2 and 3
25kg to 40kg 80mg day 1, 2 and 3
15kg to 24kg 80mg day 1 and 40mg day 2 and 3
<15kg 3mg/kg day 1 and 2mg/kg day 2 and 3

SEE
RESISTANT
CINV

DATE:
BARF SCORE:

- # review frequency after 48hr – constipation risk
- & comes as a 3 day course but may give as single dose (e.g. HD MTX) or for up to 7/7
- **ALL ARE REGULAR ONLY UNTIL 24 HOURS POST CHEMOTHERAPY THEN REVIEW**