

Area	Unfamiliar procedure	High acuity patients (excludes wards with IOA)
NICU	Exchange Transfusion Balloon septostomy Gut surgery	
Ward 23	High concentration electrolyte infusion Therapeutic heparin infusion Levosimendin infusion Pacing Berlin Heart	IOA patients
Ward 24a	High concentration electrolyte infusion Therapeutic heparin infusion Traction - if was on a non-home based ward	Unstable C spine
Ward 24b	High concentration electrolyte infusion Therapeutic heparin infusion Vac dressings - if was on a non-home based ward	New tracheostomy (first 30 days) Multiple infusion management on central line
Ward 25	High concentration electrolyte infusion Therapeutic heparin infusion	
Ward 26a	High concentration electrolyte infusion Desmopression infusion Rituximab infusion Plasmaphoresis DKA fluids/insulin infusion - if was on a non-home based ward Crisis fluid management for metabolic errors - if was on a non-home based ward	IOA patients New tracheostomy (first 30 days)
Ward 26b	CPAP/VPAP - if was on a non-home based ward	New tracheostomy (first 30 days) Tracheostomy CPAP
Ward 27		Dinutuximab infusion (over 5 days) Amofostine infusion Rituximab infusion PEG Asparaginase infusion Monoclonal antibody infusion Interleukin infusion Granulocyte infusion Cryoprecipitate transfusion Anti-thymocyte globulin infusions (ATG) Peripheral blood stem cell transfusion - day of transplant Bone marrow infusion - day of transplant