

Paediatric Crohn's Disease Activity Index: PCDAI

<i>Patient sticker here</i>

ITEM	POINTS
Abdominal pain <ul style="list-style-type: none"> None Mild (brief episodes, not interfering with activities) Moderate/severe (frequent or persistent, affecting with activities) 	0 5 10
Stools <ul style="list-style-type: none"> 0-1 liquid stools, no blood 2-5 liquid or up to 2 semi-formed with small blood Gross bleeding, >6 liquid stools or nocturnal diarrhoea 	0 5 10
Patient functioning, general well-being (Recall, 1 week) <ul style="list-style-type: none"> No limitation of activities well Occasional difficulties in maintaining age appropriate activities, below par Completely unformed 	0 5 10
Examination	
Weight <ul style="list-style-type: none"> Weight gain or voluntary weight loss Involuntary weight loss 1-9% Weight loss >10% 	0 5 10
Height <ul style="list-style-type: none"> < 1 channel decrease (or height velocity > - SD) >1 <2 channel decrease (or height velocity <-1SD >-2SD) > 2 channel decrease (or height velocity <-2SD) 	0 5 10
Abdomen <ul style="list-style-type: none"> No tenderness, or mass Tenderness, or mass without tenderness Tenderness, involuntary guarding, definite mass 	0 5 10
Peri-rectal disease <ul style="list-style-type: none"> None, asymptomatic tags 1-2 indolent fistula, scant drainage tenderness of abscess Active fistula, drainage, tenderness or abscess 	0 5 10

Extra-intestinal manifestations Fever > 38.5 x 3 days in week, arthritis, uveitis, erythema nodosum or pyoderma gangrenosum <ul style="list-style-type: none"> • None • One • Two 				0 5 10
Laboratory				
Hct%				
<10 years	11-14 (male)	11-19 (female)	15-19 (male)	
> 33	> 35	> 34	> 37	0
28-33	30-34	29-33	32-36	2.5
< 28	< 30	< 29	< 32	5
ESR mm/hr				
< 20				0
20 - 50				2.5
> 50				5
Albumin (g/L)				
> 35				0
31-34				5
< 30				10
TOTAL (SUM OF PCDAI) =				