

TYPE 2 DIABETES ACTION PLAN: 2025

Medication (Not Insulin Injections)

SCHOOL SETTING

Use in conjunction with Diabetes Management Plan.
This plan should be reviewed every year.

As kaitiaki (carers/guardians) of diabetes related services, it is a collective responsibility to establish an environment that facilitates a pathway for people with diabetes to navigate te ao mate huka - the world of diabetes¹.

The student is on a medication that **DOES NOT CAUSE** Hypoglycaemia (Hypo/Low Blood Glucose Levels)
If a student's Blood Glucose Level is less than or equal to <3.9 mmol/L they **DO NOT** require treatment or hypo food.

HIGH Hyperglycaemia (Hyper)

Blood Glucose level (BGL) greater than or equal to **>15.0 mmol/L** is well above target and requires additional action

SIGNS AND SYMPTOMS Increased thirst, extra toilet visits, poor concentration, irritability, tiredness
Note: Symptoms may not always be obvious

STUDENT WELL

- Encourage 1-2 glasses water per hour
- Return to usual activity
- Extra toilet visits may be required
- Re-check BGL in 2 hours
- Do not restrict students access to food

In 2 hours, if BGL still greater than equal to 15.0 mmol/L

CALL PARENT/CARER FOR ADVICE

STUDENT UNWELL

e.g. Vomiting
Contact parent/carer to collect student ASAP

Student's name:		Place photo of student here
Age:	DOB:	
Date:		
School name:		

Parent / Carer's name:	Contact phone:
Other contact name:	Contact phone:
Treating Medical team:	Contact phone:

BGL CHECKING TIMES

- Not required at school
- OR**
- Required at school

Frequency: