

Paediatric fluid management chart – oral diet



Name:

NHI:

Date:

Complete the chart each 24 hours

Time	Oral bottles Volume offered and volume taken	Bowel motion (use the stool tool)	Wet nappies (√)	Comments
Totals				

- If bowel motions are profuse and watery or overflowing the nappy (5e) contact your medical or nursing team and consider offering Pedialyte to drink
- If there is any blood in your child’s bowel motions, please bring them to hospital for review
- If your child has abdominal distension or acute pain please take them to hospital for review
- A temperature greater than 38 degrees should be rechecked within 20 to 30 minutes, if it remains high your child needs medical review
- If your child vomits more than 2 times in 8 hours please contact your medical or nursing team
- If your child has no wet nappies for longer than 8 hours/nil toilet urine for 12 hours please contact the team

The clinical team will add additional instructions for your child

Information provided is not intended to replace qualified medical or professional advice. For diagnosis, treatment and medication, you should consult your child’s medical specialist team.
Version 3 2020 Review date November 2022

Paediatric fluid management chart – ileostomy

Name:

NHI:

Date:

Complete the chart each 24 hours

Time	Ileostomy volume	Oral/tube intake, formula volume	Oral intake solid food	Pedialyte (Yes/No, volume given)	Wet nappies/toilet (✓)	Comments
Total						

- If stoma outputs are over _____ ml in 24 hours or over _____ ml in a 6 hour period, give Pedialyte to replace the volume lost and contact your medical or nursing team
- If there is any blood in your child’s stoma output or bowel motions, please bring them to hospital for review
- If your child has abdominal distension or acute pain please take them to hospital for review
- A temperature greater than 38 degrees should be rechecked within 20 to 30 minutes, if it remains high your child needs medical review
- If your child vomits more than 2 times in 8 hours please contact your medical or nursing team
- If your child has no wet nappies for longer than 8 hours/nil toilet urine 12 hours please contact the team

Clinical team will add additional instructions for your child

Information provided is not intended to replace qualified medical or professional advice. For diagnosis, treatment and medication, you should consult your child’s medical specialist team.
Version 3 2020 Review date November 2022

Paediatric fluid management chart – tube feed

Name:

NHI:

Date:

Complete the chart each 24 hours

Time	Continuous tube feed rate and hours/bolus feed volume	Bowel motion (use the stool tool)	Wet nappies (✓)	Comments (any oral intake)
Totals				

- If bowel motions are profuse watery/overflowing the nappy (5e) contact your medical or nursing team
- If there is any blood in your child’s bowel output or bowel motions, please bring them to hospital for review
- If your child has abdominal distension or acute pain please take them to hospital for review
- A temperature greater than 38 degrees should be rechecked within 20 to 30 minutes, if it remains high your child needs medical review
- If your child vomits more than 2 times in 8 hours please contact your medical or nursing team
- If your child has no wet nappies for longer than 8 hours/nil toilet urine for 12 hours please contact the team

The clinical team will add additional instructions for your child

Information provided is not intended to replace qualified medical or professional advice. For diagnosis, treatment and medication, you should consult your child’s medical specialist team.
Version 3 2020 Review date November 2022